REFERENCES

Allison, L.N., J.G. Hnath and W.G. Yoder. 1977. Manual of common diseases, parasites, and anomalies of Michigan fishes. Michigan Dept. Nat. Res., Lansing. Fish Mgmt. Rept. No. 8, 132 pp.

Anderson, J.R. 1967. Major land uses. Map Plates, pp. 157-159. in The National Atlas of the United States. U.S. Geological Survey. U.S. Govt. Print. Offc. Washington, D.C.

Angermier, P.L. 1985. Spatio-temporal patterns of foraging success for fishes in Illinois streams. Am. Midl. Nat. 114(2): 342-359.

_____. and J.R. Karr. 1986. Applying an index of biotic integrity based on stream fish communities: considerations in sampling and interpretation. N. Am. J. Fish. Mgmt. 6: 418-429.

Bailey, R.G. 1983. Delineation of ecosystem regions. Env. Mgmt. 7: 365-373.

Ballentine, R.K. and L.J. Guarrie (eds.). 1975. The integrity of water: a symposium. U.S. Environmental Protection Agency, Washington, D.C. 230 pp.

Balon, E.K. 1975. Reproductive guilds of fishes: a proposal and definition. J. Fish. Res. Bd. Can. 32: 821-864.

Baumann, P.C., W.D. Smith and W.K. Parland. 1987. Tumor frequencies and contaminant concentrations in brown bullhead from an industrialized river and a recreational lake. Trans. Am. Fish. Soc. 116(1): 79-86.

Becker, G.C. 1983. Fishes of Wisconsin. Univ. of Wisconsin Press, Madison. 1052 pp.

Berkman, H.E. and C.F. Rabeni. 1987. Effect of siltation on stream fish communities. Env. Biol. Fishes. 18(4): 285-294.

Berra, T.M. and R. Au. 1981. Incidences of teratological fishes from Cedar Fork Creek, Ohio. Ohio J. Sci. 81(5): 225.

Cairns, J. Jr. 1982. Artificial substrates. Ann Arbor Science Publishers, Inc., Ann Arbor, Michigan.

_____. 1986. Freshwater. In Proceedings of the workshop on cumulative environmental effects: a binational perspective. CEARC, Ottawa, Ontario and NRC, Washington, D.C. 175 pp.

Cummins, K.W. 1975. The ecology of running waters - theory and practice, pp. 278-293. in Proceedings: Sandusky River Basin Symposium. IJC International Reference Group on Great Lakes Pollution from Land Use Activities.

DeShon, J.D., D.O. McIntyre, J.T. Freda, C.D. Webster and J.P. Abrams. 1980. Volume VI, Biological evaluations, 305(b) report, 1980. Ohio EPA, Div. Surv. Water Qual. Stds., Columbus. 58 pp.

Doudoroff, P. 1951. Biological observations and toxicity bio-assays in the control of industrial waste disposal. Proc. 6th Industrial Waste Conf., Purdue Univ.

_____. and C.E. Warren. 1951. Biological indices of water pollution with special reference to fish populations. pp. 144-153, in Biological Problems in Water Pollution. U.S. Publ. Health Serv., Robt. A. Taft San. Eng. Cen., Cincinnati, Ohio.

Fausch, D.O., Karr, J.R. and P.R. Yant. 1984. Regional application of an index of biotic integrity based on stream fish communities. Trans. Amer. Fish. Soc. 113:39-55.

Gammon, J.R. 1973. The effect of thermal inputs on the populations of fish and macroinvertebrates in the Wabash River. Purdue Univ. Water Resources Res. Cen. Tech. Rep. 32. 106 pp.

Gammon, J.R. 1976. The fish populations of the middle 340 km of the Wabash River. Purdue Univ. Water Resources Res. Cen. Tech. Rep. 86. 73pp.

Gammon, J.R. 1980. The use of community parameters derived from electrofishing catches of river fish as indicators of environmental quality. pp. 335-363 in Seminar on water quality management trade-offs (point source vs. diffuse source pollution). EPA-905/9-80-009.

Gammon, J.R., A. Spacie, J.L. Hamelink, and R.L. Kaesler. 1981. Role of electrofishing in assessing environmental quality of the Wabash River, pp. 307-324 in J.M. Bates and C.I. Weber (eds.), Ecological assessments of effluent impacts on communitie of indigenous aquatic organisms. ASTM STP 703.

Hammond, E.H. 1970. Classes of land-surface form. Map Plates 61-63. in The National Atlas of the United States. U.S. Geological Survey. U.S. Govt. Print. Offc. Washington, D.C.

Herricks, E.E. and D.J. Schaeffer. 1985. Can we optimize biomonitoring? Env. Mgmt. 9: 487-492.

Hester, F.E. and J.S. Dendy. 1962. A multiple-plate sampler for aquatic macroinvertebrates. Trans. Am. Fish. Soc. 91: 420-421.

Hughes, R.M., J.H. Gakstatter, M.A. Shirazi, and J.M. Omernik. 1982. An approach for determining biological integrity in flowing waters, pp. 877-888. in T.B. Braun (ed.), Inplace Resource Inventories: Principles and Practices, A National Workshop. Soc. Amer. Foresters, Bethesda, Md.

Hughes, R.M., D.P. Larsen, and J.M. Omernik. 1986. Regional reference sites: a method for assessing stream pollution. Env. Mgmt. 10(5): 629-635.

Hughes, R.M., E. Rexstad, and C.E. Bond. 1987. The relationship of aquatic ecoregions, river basins, and physiographic provinces to the ichtyogeographic regions of Oregon. Copeia 1987: 423-432.

Jones, J.R., B.H. Tracy, J.L. Sebaugh, D.H. Hazelwood, and M.M. Smart. 1981. Biotic index tested for ability to assess water quality of Missouri Ozark streams. Trans. Am. Fish. Soc. 110(5): 627-637.

Judy, R.D., Jr., P.N. Seely, T.M. Murray, S.C. Svirsky, M.R. Whitworth, and L.S. Ischinger. 1984. 1982 National Fisheries survey. Vol. 1. Tech. Rept. Initial Findings. U.S. Fish Wildl. Serv. FWS/OBS-84/06.

Karr, J.R. 1981. Assessment of biotic integrity using fish communities. Fisheries 6 (6):21-27.

Karr, J.R. and D.R. Dudley. 1981. Ecological perspective on water quality goals. Env. Mgmt. 5(1): 55-68.

Karr, J.R., K.D. Fausch, P.L. Angermier, P.R. Yant, and I.J. Schlosser. 1986. Assessing biological integrity in running waters: a method and its rationale. Ill. Nat. Hist. Surv. Spec. Publ. 5. 28 pp.

Karr, J.R., P.R. Yant, K.D. Fausch, and I.J. Schlosser. 1987. Spatial and temporal variability of the Index of Biotic Integrity in three midwestern streams. Trans. Amer. Fish. Soc. 116(1): 1-11.

Kirsch, P.H. 1895. A report upon investigations in the Maumee River during the summer of 1893. Bull. U.S. Fish Comm. 16: 315-337.

Kuchler, A.W. 1970. Potential natural vegetation. Map plates 89-91. in The National Atlas of the United States. U.S. Geological Survey. U.S. Govt. Print. Offc. Washington, D.C.

Kuehne, R.A. and R.W. Barbour. 1983. The American darters. Univ. Press of Kentucky, Lexington. 177 pp.

Larsen, D.P., J.M. Omernik, R.M. Hughes, C.M. Rohm, T.R. Whittier, A.J. Kinney, A.L. Gallant, and D.R. Dudley. 1986. Correspondence between spatial patterns in fish assemblages in Ohio streams and aquatic ecoregions. Env. Mgmt. 10(6): 815-828.

Larsen, D.P. and D.R. Dudley. 1987. An approach for assessing attainable water quality: Ohio as a case study. Unpublished manuscript. 25 pp.

Leonard, P.M. and D.J. Orth. 1986. Application and testing of an Index of Biotic Integrity in small, cool water streams. Trans. Am. Fish. Soc. 115: 401-414.

Meek, S.E. 1889. Notes on a collection of fishes from the Maumee valley, Ohio. Proc. U.S. Nat. Mus. 2(1888): 435-440.

Mills, H.B., W.C. Starrett, and F.C. Bellrose. 1966. Man's effect on the fish and wildlife of the Illinois River. Ill. Nat. Hist. Surv. Biol. Notes 57. 27 pp.

Novotny D.W. and G.R. Priegel. 1974. Electrofishing boats, improved designs, and operational guidelines to increase the effectiveness of boom shockers. Wisc. DNR Tech. Bull. No. 73, Madison WI. 48 pp.

Odum, E.P., J.T. Finn, and E.H. Franz. 1979. Perturbation theory and the subsidy-stress gradient. BioScience 29: 349-352.

Ohio Department of Natural Resources. 1960. Gazetteer of Ohio streams. Ohio DNR, Div. of Water, Columbus, Ohio. Ohio Water Plan Inventory Rept. No. 12. 179 p.

_____. 1967. Drainage areas of Ohio streams. Supplement to gazetteer of Ohio streams. Ohio Water Plan Inv. Rept. 12a. 61 pp.

Ohio Environmental Protection Agency. 1987a. Biological criteria for the protection of aquatic life: Volume III. Standardized field and laboratory methods for assessing fish and macroinvertebrate communities. Division of Water Quality Monitoring and Assessment, Surface Water Section, Columbus, Ohio.

_____. 1987b. Biological criteria for the protection of aquatic life: Volume I. The role of biological data in water quality assessment. Division of Water Quality Monitoring and Assessment, Surface Water Section, Columbus, Ohio.

_____. 1986. The cost of biological field monitoring. Division of Water Quality Monitoring and Assessment, Columbus, Ohio. 6 pp. (mimeo)

_____. 1984. Implementation manual for water quality standards. Div. Water Qual. Monitoring and Assess., Eval. and Stds. Sect., Columbus.

_____. 1983a. Biological and water quality study of Big Darby Creek, Union and Madison Counties, Ohio. C. Yoder (ed.). Div. Wastewater Poll. Contr., Surveillance and Stds. Section, Columbus.

_____. 1983b. The PEMSO system: stream network file users manual, report no. 4. Offc. Planning Coord., Columbus. 38 pp.

_____. 1982. Biological and water quality study of the lower mainstem of the Great Miami River. C.O. Yoder (ed.). Ohio EPA Tech. Rept. 82/12. Div. Wastewater Poll. Contr., Columbus. 219 pp.

Omernik, J.M. 1987. Ecoregions of the conterminous United States. Ann. Assoc. Amer. Geogr. 77(1): 118-125.

Post, G. 1983. Textbook of fish health. TFH Publications, Inc., Neptune City. 256 pp.

Smith, P.W. 1968. An assessment of changes in the fish fauna of two Illinois rivers and its bearing on their future. Trans. Ill. State Acad. Sci. 61(1): 31-45.

_____. 1971. Illinois streams: a classification based on their fishes and an analysis of factors responsible for the disappearance of native species. Ill. Nat. Hist. Surv. Biol. Notes 76.

_____. 1979. The fishes of Illinois. Univ. Illinois Press, Urbana. 314 pp.

Thoma, R.T., E.T. Rankin, M. Smith, and R. Sanders. 1987. An objective method for ranking the general intolerance of stream fishes. Ohio Fish and Wildlife Conference, Columbus, Ohio (poster session).

Trautman, M.B. 1939. The effects of man-made modifications on the fish fauna in Lost and Gordon Creeks, Ohio, between 1887-1938. Ohio J. Sci. 39(5): 275-288.

_____. 1942. Fish distribution and abundance correlated with stream gradient as a consideration in stocking programs. Trans. 7th N. Am. Wildl. Conf. 7: 211-224.

_____. and R.K. Gartman. 1974. Re-evaluation of the effects of man-made modifications of Gordon Creek between 1887 and 1973 and especially as regards its fish fauna. Ohio J. Sci. 74(3): 162-173.

_____. 1981. The fishes of Ohio. (2nd edition). Ohio State Univ. Press, Columbus. 782 p.

Vannote, R.L., G.W. Minshall, K.W. Cummins, J.R. Sedell and C.K. Cushing. 1980. The river continuum concept. Can. J. Fish. Aquat. Sci. 37:130-137.

Vincent, R. 1971. River electrofishing and fish population estimates. Prog. Fish Cult. 33(3): 163-169.

WAPORA. 1978. Fish populations and water quality of the lower 200 miles of the West Fork and mainstem White River, Indiana. WAPORA, Inc., Cincinnati, Ohio. 47 pp.

Whittier, T.R., D.P. Larsen, R.M. Hughes, C.M. Rohm, A.L. Gallant, and J.M. Omernik. 1987. The Ohio stream regionalization project: a compendium of results. U.S. EPA - Freshwater Res. Lab, Corvallis, OR. EPA/600/3-87/025. 163 pp.

Yoder, C.O., P. Albeit, and M.A. Smith. 1981. The distribution and abundance of fishes in the mainstem Scioto River as affected by pollutant loadings. Ohio EPA Tech. Rept. 81/3. Columbus. 118 pp.